

POINTWISE BOUNDED ASYMPTOTIC MORPHISMS AND THOMSEN'S NON-STABLE k -THEORY

E.J. Beggs

Dept of Mathematics
University of Wales Swansea
SA2 8PP U.K.

E.J.Beggs@swansea.ac.uk

Abstract

In this paper I show that pointwise bounded asymptotic morphisms between separable metrisable locally convex $*$ -algebras induce continuous maps between the quasi-unitary groups of the algebras, provided that the algebras support a certain amount of functional calculus. This links the asymptotic morphisms directly to Thomsen's non-stable definition of k -theory in the C^* algebra case. A result on composition of asymptotic morphisms is also given.

1 Introduction

Thomsen defined a non-stable version of K -theory, called k -theory, in terms of the topology of the unitary group of a C^* algebra [6]. Here we link the pointwise bounded asymptotic morphisms (PBAMs) on separable metrisable locally convex (SMLC) algebras to the topology of their unitary groups (for $*$ -algebras). For other algebras, it may be more appropriate to consider the group of invertible elements, for which a similar approach could be taken, but in this paper we concentrate on $*$ -algebras and the unitary group.

To avoid problems with algebras which do not contain a unit, Thomsen used the following procedure, following Palmer [4, 5]. On an algebra A we define an associative binary operation $a \bullet b = a + b + ab$, which has 2-sided unit $0 \in A$. Note that if we had a unit 1 for algebra multiplication, we would have $(1 + a)(1 + b) = 1 + a \bullet b$. By this means it can be seen that this

construction is really standard, though phrased in slightly unusual terms. We define $\text{gl}(A)$, the quasi-invertibles in an algebra A , and $\mathcal{U}(A)$, the quasi-unitaries, by

$$\text{gl}(A) = \{a \in A : \exists a' \in A \ a \bullet a' = a' \bullet a = 0\} , \ \mathcal{U}(A) = \{u \in A : u \bullet u^* = u^* \bullet u = 0\} .$$

It is easily seen that both of these are groups under the $\bullet$ operation, with identity 0. Thomsen's non-stable k -groups for a C^* algebra E are defined as $k_n(E) = \pi_{n+1}(\mathcal{U}(E))$.

The idea of asymptotic morphisms for C^* algebras was introduced by Connes and Higson [2] for E -theory, and later considered in the non-stable C^* case by Dădărlat [3]. In [1] PBAMs were introduced to generalise the idea of asymptotic morphism to SMLC algebras, and examples were given. The definition of a PBAM will be given later in this paper. The general idea of all these constructions is to have a family of maps $f_t : A \rightarrow B$ between algebras indexed by $t \in [0, \infty)$, where f_t becomes more like an algebra map as $t \rightarrow \infty$.

Under a certain assumption about functional calculus on SMLC $*$ -algebras A and B , we show that a PBAM $f : A \times [0, \infty) \rightarrow B$ induces a unique topological homotopy class $\mathcal{U}(f) \in [\mathcal{U}(A), \mathcal{U}(B)]$. In the C^* algebra case, this gives a map $f_* : k_*(A) \rightarrow k_*(B)$ by composition, $f_*[h] = [\mathcal{U}(f) \circ h]$. The question naturally arises of whether $\mathcal{U}$ is a functor from SMLC $*$ -algebras and PBAMs to topological spaces. The answer is no, as SMLC $*$ -algebras and PBAMs themselves do not form a category. However there is a compatibility condition between PBAMs $f : A \times [0, \infty) \rightarrow B$ and $g : B \times [0, \infty) \rightarrow C$ which ensures that there is a composition $g \circ_\phi f : A \times [0, \infty) \rightarrow C$, defined up to homotopy of PBAMs. We shall see that this same condition also implies that $\mathcal{U}(g \circ_\phi f) = \mathcal{U}(g) \circ \mathcal{U}(f) \in [\mathcal{U}(A), \mathcal{U}(C)]$. I choose to retain the compatibility condition rather than restrict to a category where composition is automatic (and there are such categories) in order not to unnecessarily restrict the results.

2 Asymptotic results and definitions on metric spaces

Lemma 2.1 *Let X be a separable metric space. Suppose that there is a function $\Psi : X \times X \times [0, \infty) \times [0, \infty) \rightarrow \{\text{true}, \text{false}\}$ with the following property: For all $x \in X$ there is a $\delta(x) > 0$ and a $Q(x) \geq 0$ so that for all $q \geq Q$ there is a $R(x, q) \geq 0$ and an $\epsilon(x, q) > 0$ for which*

$$\forall y \in X \ \forall t \geq Q \ \forall s \geq R \quad [t \in (q - \epsilon, q + \epsilon) \text{ and } d(x, y) < \delta] \Rightarrow \Psi(x, y, s, t) .$$

Then there are continuous functions $\alpha : X \rightarrow [0, \infty)$ and $\phi : [0, \infty) \rightarrow [0, \infty)$ (increasing) so that

$$\forall y \in X \quad \exists x \in X \quad \forall t \geq \alpha(y) \quad \forall s \geq \phi(t) \quad [d(x, y) < \delta(x) \text{ and } \Psi(x, y, s, t)] .$$

Proof Part 1: Given $x \in X$ and $r \geq Q(x)$, there is an open cover $(q - \epsilon(x, q), q + \epsilon(x, q))$ of the interval $[Q, r]$ indexed by $q \in [Q, r]$. By compactness of $[Q, r]$ there is a finite subcover indexed by $q_1, \dots, q_n$. Let $S(x, r)$ be the maximum of the $R(x, q_i)$. Then

$$\forall y \in X \quad \forall t \in [Q, r] \quad \forall s \geq S \quad d(x, y) < \delta \Rightarrow \Psi(x, y, s, t) .$$

Part 2: Take a cover of X by open balls center $x \in X$ radius $\delta(x)$. There is a countable subcover of X by open balls B_i (integer $i \geq 0$) center x_i radius $\delta(x_i)$. Now define an increasing sequence $\phi_i \in [0, \infty)$ by using the recursive inequalities

$$\forall j \in \{0, \dots, i\} \quad \text{if } i \geq Q(x_j) \text{ then } \phi_i \geq S(x_j, i + 1) .$$

Construct the graph of the continuous increasing function $\phi : [0, \infty) \rightarrow [0, \infty)$ by joining the dots $(0, \phi_0), (1, \phi_1), \dots$ with straight lines. Take a continuous partition of unity $\theta_j : X \rightarrow [0, 1]$ with locally finite supports, where θ_j has support in B_j , and define

$$\alpha(x) = \sum_j (1 + \max\{Q(x_j), j\}) \theta_j(x) .$$

Part 3: For a given $y \in X$ we choose the minimum value of $\max\{Q(x_j), j\}$ over the finite number of indices j for which $\theta_j(y) \neq 0$. If this minimum is $\max\{Q(x_k), k\}$ where $\theta_k(y) \neq 0$, it follows that $\alpha(y) \geq 1 + \max\{Q(x_k), k\}$ and $y \in B_k$. Now if $t \geq 1 + \max\{Q(x_k), k\}$ then there is an $i \in \mathbb{N}$ with $i + 1 \geq t \geq i \geq Q(x_k)$ and $i \geq k$. Now if $s \geq \phi(t)$ then $s \geq \phi_i$, so $s \geq S(x_k, i + 1)$. It follows that $\Psi(x_k, y, s, t)$. $\square$

Lemma 2.2 For $n \in \mathbb{N}$, let X_n be a separable metric space. Suppose that there are functions $\Psi_n : X_n \times X_n \times [0, \infty) \times [0, \infty) \rightarrow \{\text{true}, \text{false}\}$ (for $n \in \mathbb{N}$) which satisfy the property in the statement of lemma 2.1, using functions δ_n, Q_n, R_n and ϵ_n . Then there is a continuous increasing function $\psi : [0, \infty) \rightarrow [0, \infty)$ and continuous functions $\beta_n : X_n \rightarrow [0, \infty)$ so that

$$\forall n \in \mathbb{N} \quad \forall y \in X_n \quad \exists x \in X_n \quad \forall t \geq \beta_n(y) \quad \forall s \geq \psi(t) \quad [d(x, y) < \delta_n(x) \text{ and } \Psi_n(x, y, s, t)] .$$

Proof By using lemma 2.1 on each of the Ψ_n , we get continuous functions $\alpha_n : X_n \rightarrow [0, \infty)$ and $\phi_n : [0, \infty) \rightarrow [0, \infty)$. Now we make a new continuous increasing function $\psi : [0, \infty) \rightarrow [0, \infty)$ which satisfies the condition that if $t \geq n$ then $\psi(t) \geq \phi_n(t)$. Now set $\beta_n(x) = \max\{n, \alpha_n(x)\}$. $\square$

Definition 2.3 For metric spaces X and Y , the function $F : X \times [0, \infty) \rightarrow Y$ is said to be strongly asymptotically continuous if, for all $\epsilon > 0$ and all $x \in X$ there is an $\eta(x, \epsilon) > 0$ and a $P(x, \epsilon) \geq 0$ so that for all $x' \in X$ with $d(x, x') < \eta$ we have $d(F_t(x), F_t(x')) < \epsilon$ for all $t \geq P$.

Lemma 2.4 For metric spaces X and Y , suppose that the function $F : X \times [0, \infty) \rightarrow Y$ is continuous and strongly asymptotically continuous. Then, given $\epsilon > 0$ and $x \in X$, there are $\bar{\eta}(x, \epsilon) > 0$ and $\bar{P}(x, \epsilon) \geq 0$ so that, given $q \geq \bar{P}$ there is a $\bar{\delta}(x, \epsilon, q) > 0$ so that for all $x' \in X$ and all $t \geq 0$,

$$[|t - q| < \bar{\delta} \text{ and } d(x, x') < \bar{\eta}] \Rightarrow d(F_q(x), F_t(x')) < \epsilon .$$

Proof From definition 2.3 we take $\bar{\eta} = \eta(x, \epsilon/2) > 0$ and $P(x, \epsilon/2) \geq 0$, and set $\bar{P} = P + 1$. By continuity of F , given $q \geq \bar{P}$ there is a $1 > \bar{\delta}(x, \epsilon, q) > 0$ so that $|t - q| < \bar{\delta}$ implies $d(F_q(x), F_t(x)) < \epsilon/2$. As $|t - q| < \bar{\delta}$ implies $t \geq P$, we have $d(F_t(x), F_t(x')) < \epsilon/2$ for all $x' \in X$ with $d(x, x') < \bar{\eta}$. $\square$

3 Some functional calculus

Definition 3.1 Let A be a topological $*$ -algebra, and take A_{sad} to be the subset of self adjoint elements, i.e. those $a \in A$ for which $a^* = a$. A is said to have the inverse square root property if there is a convex open set $V_A \subset A_{sad}$ containing $0 \in A$ and a continuous function $\vartheta : V_A \rightarrow A_{sad}$ so that

- 1) $a \bullet \vartheta(a) = \vartheta(a) \bullet a$ for all $a \in V_A$,
- 2) $a \bullet (\vartheta(a) \bullet \vartheta(a)) = 0$ for all $a \in V_A$,
- 3) for all $a \in V_A$, a is $\bullet$ -invertible,
- 4) $\vartheta(0) = 0$.

Lemma 3.2 If $a \in A$ has $a^* \bullet a \in V_A$ and $a \bullet a^* \in V_A$, then $a \in \text{gl}(A)$ and $a \bullet \vartheta(a^* \bullet a) \in \mathcal{U}(A)$.

Proof Set $u = a \bullet \vartheta(a^* \bullet a)$. Then

$$u^* \bullet u = \vartheta(a^* \bullet a) \bullet (a^* \bullet a) \bullet \vartheta(a^* \bullet a) = (a^* \bullet a) \bullet (\vartheta(a^* \bullet a) \bullet \vartheta(a^* \bullet a)) = 0 .$$

Since $a^* \bullet a$ and $a \bullet a^*$ are $\bullet$ -invertible, there are $b \in A$ and $c \in A$ so that

$$a^* \bullet a \bullet b = b \bullet a^* \bullet a = 0 \quad \text{and} \quad a \bullet a^* \bullet c = c \bullet a \bullet a^* = 0 .$$

Now

$$(a^* \bullet a) \bullet (b \bullet a^*) = 0 \bullet a^* = a^* = a^* \bullet 0 = a^* \bullet (a \bullet a^* \bullet c) = (a^* \bullet a) \bullet (a^* \bullet c) ,$$

so since $a^* \bullet a$ is $\bullet$ -invertible we deduce that $b \bullet a^* = a^* \bullet c$ is the $\bullet$ -inverse of a . The formula

$$((a^* \bullet a) \bullet \vartheta(a^* \bullet a)) \bullet \vartheta(a^* \bullet a) = 0$$

shows that $\vartheta(a^* \bullet a)$ is $\bullet$ -invertible, so u is $\bullet$ -invertible, and so $u \bullet u^* = 0$, i.e. $u \in \mathcal{U}(A)$. $\square$

Example 3.3 Any Banach $*$ -algebra A has the inverse square root property. This can be seen by taking $V_A = \{a \in A_{\text{sad}} : |a| < \frac{1}{2}\}$, and using functional calculus with the function $\vartheta(x) = (1+x)^{-1/2} - 1$. The easiest way to see that this works is to observe that ϑ has a Taylor series with radius of convergence 1, and the fact that there is no constant term in the series means that we do not require A to have a unit.

Example 3.4 The algebra $C^\infty(M, \mathbb{C})$ of smooth complex valued functions on a compact manifold M (with the smooth topology) has the inverse square root property. We set $V = \{f \in C^\infty(M, \mathbb{R}) : \forall m \in M \ |f(m)| < \frac{1}{2}\}$, and $\vartheta(f)(m) = (1 + f(m))^{-1/2} - 1$.

4 Pointwise bounded asymptotic morphisms

Suppose that both A and B are separable metrisable locally convex $*$ -algebras (SMLC $*$ -algebras for short). Recall that a metrisable locally convex topological vector space has topology defined by a countable number of seminorms, which we write $|\cdot|_0, |\cdot|_1$, etc. We can change any of the seminorms to a stronger continuous seminorm without altering the topology. It will be convenient to use this fact to assume that the seminorms are non-decreasing, i.e. that $|b|_n \leq |b|_{n+1}$ for all $n \in \mathbb{N}$ and $b \in B$. In a SMLC $*$ -algebra B we assume that the multiplication and

star operations are continuous. It will also be convenient to assume, by strengthening seminorms where appropriate, that for all $b, b' \in B$ and $n \in \mathbb{N}$, $|b^*|_n \leq |b|_{n+1}$ and $|bb'|_n \leq |b|_{n+1}|b'|_{n+1}$. This implies the following result for $b, c \in B$, which is used later:

$$|b^* \bullet b - c^* \bullet c|_n \leq 2 |b - c|_{n+2} (1 + |b|_{n+2} + |b - c|_{n+2}) . \quad (1)$$

Definition 4.1 *The map $f : A \times [0, \infty) \rightarrow B$ is called pointwise asymptotically bounded if, for all $a \in A$, the set $\{f_t(a) : t \geq 0\}$ is bounded in B . In effect this means that for every $n \in \mathbb{N}$ the subset of the reals $\{|f_t(a)|_n : t \geq 0\}$ is bounded above.*

Definition 4.2 *If A and B are SMLC $*$ -algebras, call a continuous function $f : A \times [0, \infty) \rightarrow B$ a pointwise bounded asymptotic morphism (PBAM for short) if*

- a) $f : A \times [0, \infty) \rightarrow B$ is pointwise asymptotically bounded (see 4.1),
- b) f is strongly asymptotically continuous (see 2.3),
- c) $f_t(a)^* - f_t(a^*) \rightarrow 0$ as $t \rightarrow \infty$ for all $a \in A$,
- d) $\lambda f_t(a) - f_t(\lambda a) \rightarrow 0$ as $t \rightarrow \infty$ for all $a \in A$ and all $\lambda \in \mathbb{C}$,
- e) $f_t(a) + f_t(a') - f_t(a + a') \rightarrow 0$ as $t \rightarrow \infty$ for all $a, a' \in A$,
- f) $f_t(a) f_t(a') - f_t(a a') \rightarrow 0$ as $t \rightarrow \infty$ for all $a, a' \in A$.

Note that the additive condition (e) implies that $f_t(0) \rightarrow 0$ as $t \rightarrow \infty$.

5 Pointwise bounded asymptotic morphisms and quasi-unitary groups

In this section we suppose that A and B are SMLC $*$ -algebras, that B has the inverse square root property (see 3.1), and that $f : A \times [0, \infty) \rightarrow B$ is a PBAM.

Lemma 5.1 *For every $u \in \mathcal{U}(A)$ there is a $T(u) \geq 0$ and an open neighbourhood $O(u)$ of u in A so that*

$$\forall t \geq T(u) \quad \forall v \in O(u) \quad f_t(v) \bullet f_t(v)^* \in V_B \text{ and } f_t(v)^* \bullet f_t(v) \in V_B .$$

Proof First note that, for the given $u \in \mathcal{U}(A)$, $f_t(u) \bullet f_t(u)^* \rightarrow 0$ and $f_t(u)^* \bullet f_t(u) \rightarrow 0$ as $t \rightarrow \infty$. To see this, apply f_t to $u^* \bullet u = 0$ to get $f_t(u) + f_t(u^*) + f_t(u^*) f_t(u) \rightarrow 0$ as $t \rightarrow \infty$. Now we use $f_t(u)^* - f_t(u^*) \rightarrow 0$ as $t \rightarrow \infty$ together with the fact that $f_t(u)$ is bounded for $t \geq 0$ to show that $f_t(u)^* \bullet f_t(u) \rightarrow 0$. Similarly for $f_t(u) \bullet f_t(u)^* \rightarrow 0$.

As V_B is an open neighbourhood of 0 in B_{sad} , there is an $n \geq 0$ and an $\epsilon > 0$ so that $\{b \in B_{sad} : |b|_n < 2\epsilon\} \subset V_B$. As f is pointwise bounded, there is a $K \geq 0$ so that $|f_t(u)|_{n+2} \leq K$ for all t . Choose $\delta > 0$ so that $2\delta(1 + K + \delta) < \epsilon$. By strong asymptotic continuity there is a neighbourhood $O(u)$ of u in A and a $P \geq 0$ so that $|f_t(v) - f_t(u)|_{n+2} \leq \delta$ for all $t \geq P(u)$ and all $v \in O(u)$. Then by (1), $|f_t(v)^* \bullet f_t(v) - f_t(u)^* \bullet f_t(u)|_n < \epsilon$. Likewise we get $|f_t(v) \bullet f_t(v)^* - f_t(u) \bullet f_t(u)^*|_n < \epsilon$. Finally choose $T \geq P$ so that $|f_t(u)^* \bullet f_t(u)|_n < \epsilon$ and $|f_t(u) \bullet f_t(u)^*|_n < \epsilon$ for all $t \geq T$. $\square$

Lemma 5.2 *There is a continuous function $\alpha : \mathcal{U}(A) \rightarrow [0, \infty)$ so that*

$$\forall v \in \mathcal{U}(A) \quad \forall t \geq \alpha(v) \quad f_t(v) \bullet f_t(v)^* \in V_B \text{ and } f_t(v)^* \bullet f_t(v) \in V_B .$$

Proof From lemma 5.1 take a countable subcover $O(u_n)$ ($n \in \mathbb{N}$) of $\mathcal{U}(A)$, and a locally finite continuous partition of unity θ_n on $\mathcal{U}(A)$, with the support of θ_n contained in $O(u_n)$. Now set $\alpha = \sum T(u_n) \theta_n$. $\square$

Proposition 5.3 *There is a well defined homotopy class $\mathcal{U}(f) \in [\mathcal{U}(A), \mathcal{U}(B)]$ with representative function $\tilde{f}_\alpha(v) = f_{\alpha(v)}(v) \bullet \vartheta(f_{\alpha(v)}(v)^* \bullet f_{\alpha(v)}(v))$, where α is the function in lemma 5.2. Here well defined means independent of the choice of α .*

Proof Note that the function $\tilde{f}_\alpha$ takes values in $\mathcal{U}(B)$ by lemma 3.2. Suppose that we have $\gamma : \mathcal{U}(A) \rightarrow [0, \infty)$ with $\gamma(u) \geq \alpha(u)$ for all $u \in \mathcal{U}(A)$. Then there is a homotopy $H : \mathcal{U}(A) \times [0, 1] \rightarrow \mathcal{U}(B)$ connecting $\tilde{f}_\alpha$ and $\tilde{f}_\gamma$ defined by $H(v, p) = \tilde{f}_{p\alpha + (1-p)\gamma}(v)$. Now if β is another choice of function in lemma 5.2, we see that $\tilde{f}_\alpha$ and $\tilde{f}_\beta$ are homotopic, as they are both homotopic to $\tilde{f}_{\max\{\alpha, \beta\}}$. $\square$

6 Homotopy invariance

In this section we suppose that A and B are SMLC $*$ -algebras, and that f and $g : A \times [0, \infty) \rightarrow B$ are two PBAMs.

Definition 6.1 *The PBAMs f and g are said to be pointwise bounded asymptotically homotopy equivalent (PBA homotopic for short) if there is a PBAM $h : A \times [0, \infty) \rightarrow C([0, 1], B)$ so that $h_t(a)(0) = f_t(a)$ and $h_t(a)(1) = g_t(a)$. (Use the supremum metric on $C([0, 1], B)$, the continuous functions from $[0, 1]$ to B .)*

Proposition 6.2 *If f and g are PBA homotopic then $\mathcal{U}(f) = \mathcal{U}(g) \in [\mathcal{U}(A), \mathcal{U}(B)]$.*

Proof Take the homotopy to be $h : A \times [0, \infty) \rightarrow C([0, 1], B)$. By lemma 5.2 there are functions α, β and $\gamma : \mathcal{U}(A) \rightarrow [0, \infty)$ corresponding to f, g and h respectively. Define $\eta : \mathcal{U}(A) \rightarrow [0, \infty)$ by $\eta(u) = \max\{\alpha(u), \beta(u), \gamma(u)\}$. Now $\tilde{f}_\alpha$ is homotopic to $\tilde{f}_\eta$, and $\tilde{g}_\beta$ is homotopic to $\tilde{g}_\eta$. Finally $\tilde{f}_\eta$ and $\tilde{g}_\eta$ are homotopic by the homotopy $H : \mathcal{U}(A) \times [0, 1] \rightarrow \mathcal{U}(B)$ defined by $H(v, p) = \tilde{h}_\eta(v)(p)$, using the fact that $\mathcal{U}(C([0, 1], B)) = C([0, 1], \mathcal{U}(B))$. $\square$

7 Composition of pointwise bounded asymptotic morphisms

In this section we suppose that A, B and C are SMLC *-algebras, and that $f : A \times [0, \infty) \rightarrow B$ and $g : B \times [0, \infty) \rightarrow C$ are PBAMs. We will show that under certain conditions it is possible to compose f and g . To save a lot of time we will use the results on composing strongly asymptotic morphisms in [1].

Definition 7.1 *For PBAMs f and g , and a continuous increasing function $\phi : [0, \infty) \rightarrow [0, \infty)$ we define $g \circ_\phi f : A \times [0, \infty) \rightarrow C$ by $(g \circ_\phi f)_t(a) = g_{\phi(t)}(f_t(a))$. The function ϕ is called a valid reparameterisation for f and g if*

- a) $g \circ_\phi f$ is a PBAM, and
- b) for every continuous increasing function $\theta : [0, \infty) \rightarrow [0, \infty)$ with $\theta(t) \geq \phi(t) \forall t \in [0, \infty)$, $g \circ_\theta f$ is also a PBAM, and is PBA homotopic to $g \circ_\phi f$, using $h : A \times [0, \infty) \rightarrow C([0, 1], B)$ given by $h_t(a)(p) = g_{p\phi(t)+(1-p)\theta(t)}(f_t(a))$.

Definition 7.2 *The PBAMs f and g are said to satisfy conditions C1 and C3 if:*

C1): For all $a \in A$ and all $\nu > 0$ there is a $\xi(a, \nu) > 0$ and a $Q'(a, \nu) \geq 0$ so that, for all $t \geq Q'$ there is a $S'(t, a, \nu) \geq 0$ so that

$$[d(f_t(a), b) < \xi \text{ and } s \geq S'] \Rightarrow d(g_s(f_t(a)), g_s(b)) < \nu.$$

C3): For all $a \in A$ and all $n \in \mathbb{N}$ there is a $Q_n(a) \geq 0$ and an $M_n(a) \geq 0$ so that, for all $t \geq Q_n(a)$ there is a $S_n(a, t) \geq 0$ so that for $s \geq S_n$ we have $|g_s(f_t(a))|_n \leq M_n$.

Lemma 7.3 *If the PBAMs f and g satisfy conditions C1 and C3, then the property $\Psi_n : A \times A \times [0, \infty) \times [0, \infty) \rightarrow \{\text{true}, \text{false}\}$ defined by $\Psi_n(x, y, s, t)$ being the logical value of $|g_s(f_t(y))|_n \leq M_n(x) + 1$ satisfies the conditions of lemma 2.1.*

Proof Choose $\nu > 0$ so that for the translation invariant metric d on C , $d(c, c') < \nu$ implies $|c - c'|_n < 1$. By applying C3 to $x \in A$, we get $Q_n(x) \geq 0$ and an $M_n(x) \geq 0$ so that, for all $q \geq Q_n(x)$ there is a $S_n(x, q) \geq 0$ so that for $s \geq S_n$ we have $|g_s(f_q(x))|_n \leq M_n$. By applying C1, we get $\xi(x, \nu) > 0$ and a $Q'(x, \nu) \geq 0$ so that, for all $q \geq Q'$ there is a $S'(q, x, \nu) \geq 0$ so that

$$[d(f_q(x), b) < \xi \text{ and } s \geq S'] \Rightarrow |g_s(f_q(x)) - g_s(b)|_n < 1 .$$

By lemma 2.4 there are $\bar{\eta}(x, \xi) > 0$ and $\bar{P}(x, \xi) \geq 0$ so that, for $q \geq \bar{P}$ there is a $\bar{\delta}(x, \xi, q) > 0$ so that for all $y \in A$ and all $t \geq 0$,

$$[d(x, y) < \bar{\eta} \text{ and } |t - q| < \bar{\delta}] \Rightarrow d(f_q(x), f_t(y)) < \xi .$$

Now set $\delta(x) = \bar{\eta}(x, \nu)$ and $Q(x) = \max\{Q_n(x), Q'(x, \nu), \bar{P}(x, \xi)\}$. Given $q \geq Q$ set $\epsilon(x, q) = \bar{\delta}(x, \xi, q)$ and $R(x, q) = \max\{S'(q, x, \nu), S_n(x, q)\}$. $\square$

Theorem 7.4 *If the PBAMs f and g satisfy conditions C1 and C3, then there is a continuous increasing function $\phi : [0, \infty) \rightarrow [0, \infty)$ which is a valid reparameterisation for f and g .*

Proof First note that (C1 and C3) implies C2 from [1] for the $*$ -operation, scalar multiplication, addition and algebra multiplication. To see this we use the inequalities

$$\begin{aligned} |g_s(f_t(a))^* - g_s(b)^*|_n &\leq |g_s(f_t(a)) - g_s(b)|_{n+1} , \\ |\lambda g_s(f_t(a)) - \mu g_s(b)|_n &\leq |\lambda - \mu| |g_s(f_t(a))|_n + \\ &\quad (|\lambda| + |\lambda - \mu|) |g_s(f_t(a)) - g_s(b)|_n , \\ |g_s(f_t(a)) + g_s(f_t(a')) - g_s(b) - g_s(b')|_n &\leq |g_s(f_t(a)) - g_s(b)|_n + |g_s(f_t(a')) - g_s(b')|_n , \\ |g_s(f_t(a)) g_s(f_t(a')) - g_s(b) g_s(b')|_n &\leq |g_s(f_t(a))|_{n+1} |g_s(f_t(a')) - g_s(b')|_{n+1} + \\ &\quad |g_s(f_t(a)) - g_s(b)|_{n+1} \times \end{aligned}$$

$$(|g_s(f_t(a'))|_{n+1} + |g_s(f_t(a')) - g_s(b')|_{n+1}) . \quad (2)$$

From [1] we now know that there is a valid reparameterisation (for strongly asymptotic morphisms) $\psi : [0, \infty) \rightarrow [0, \infty)$ so that $g \circ_\psi f$ is a **strongly** asymptotic morphism. From 7.3 and 2.2 there are functions $\beta_n : A \rightarrow [0, \infty)$ ($n \in \mathbb{N}$) and $\theta : [0, \infty) \rightarrow [0, \infty)$ so that, given $n \in \mathbb{N}$ and $y \in A$, there is an $x \in A$ so that

$$t \geq \beta_n(y) \text{ and } s \geq \theta(t) \Rightarrow |g_s(f_t(y))|_n \leq M_n(x) + 1 .$$

Let $\phi(y) = \max\{\psi(y), \theta(y)\}$. Then $g \circ_\phi f$ is a PBAM, and if $\chi : [0, \infty) \rightarrow [0, \infty)$ has $\chi(t) \geq \phi(t)$ $\forall t \in [0, \infty)$, then the map $h : A \times [0, \infty) \rightarrow C([0, 1], B)$ given by $h_t(a)(p) = g_{p\phi(t) + (1-p)\chi(t)}(f_t(a))$ is a PBA homotopy between $g \circ_\phi f$ and $g \circ_\chi f$. $\square$

8 A conditional functoriality

Suppose that A , B and C are SMLC $*$ -algebras, and that B and C have the inverse square root property. We are given PBAMs $f : A \times [0, \infty) \rightarrow B$ and $g : B \times [0, \infty) \rightarrow C$ which satisfy $C1$ and $C3$.

Lemma 8.1 *Given $x \in \mathcal{U}(A)$, there is an $N(x) \geq 0$ and a $\nu(x) > 0$ so that, for $q \geq N$ there is an $L(x, q) \geq 0$ so that*

$$\forall z \in B \quad [d(f_q(x), z) < \nu \text{ and } s \geq L] \Rightarrow [g_s(z)^* \bullet g_s(z) \in V_C \text{ and } g_s(z) \bullet g_s(z)^* \in V_C] .$$

Proof Part 1: We shall only consider the $g_s(z)^* \bullet g_s(z) \in V_C$ part, the other is almost identical. Choose $\epsilon > 0$ and $n \geq 0$ so that for $c \in C$, $|c|_n < \epsilon$ implies $c \in V_C$. Since $g_s(0) \rightarrow 0$ as $s \rightarrow \infty$, and using the strong asymptotic continuity of g , there is a $\eta > 0$ and $P \geq 0$ so that

$$[d(b, 0) < \eta \text{ and } s \geq P] \Rightarrow |g_s(b)|_n < \epsilon/4 . \quad (3)$$

Given $x \in \mathcal{U}(A)$, by $C3$ and (1) there is a $Q_{n+2}(x) \geq 0$ and $M_{n+2}(x) \geq 0$ so that for any $q \geq Q_{n+2}$ there is a $S_{n+2}(x, q)$ so that for $s \geq S_{n+2}$,

$$\begin{aligned} |g_s(z)^* \bullet g_s(z) - g_s(f_q(x))^* \bullet g_s(f_q(x))|_n &\leq 2|g_s(z) - g_s(f_q(x))|_{n+2} \times \\ &\quad (1 + M_{n+2} + |g_s(z) - g_s(f_q(x))|_{n+2}) . \end{aligned} \quad (4)$$

Now choose $\delta(x) > 0$ so that $2\delta(1 + M_{n+2} + \delta) < \epsilon/2$. By *C1* there is a $\xi(x)$ and a $Q'(x)$ so that, for all $q \geq Q'$ there is a $S'(x, q)$ so that

$$[d(f_q(x), z) < \xi \text{ and } s \geq S'] \Rightarrow |g_s(z) - g_s(f_q(x))|_{n+2} < \delta. \quad (5)$$

As f is a PBAM we know that $f_t(x)^* \bullet f_t(x) - f_t(x^* \bullet x) \rightarrow 0$ as $t \rightarrow \infty$, and as $x \in \mathcal{U}(A)$ this means $f_t(x)^* \bullet f_t(x) \rightarrow 0$. Choose $N'(x) \geq 0$ so that $d(f_t(x)^* \bullet f_t(x), 0) < \eta$ for all $t \geq N'$. Given $q \geq 0$, as g is a PBAM we know that $g_s(f_q(x))^* \bullet g_s(f_q(x)) - g_s(f_q(x)^* \bullet f_q(x)) \rightarrow 0$ as $s \rightarrow \infty$. Choose $S''(x, q) \geq 0$ so that

$$s \geq S'' \Rightarrow |g_s(f_q(x))^* \bullet g_s(f_q(x)) - g_s(f_q(x)^* \bullet f_q(x))|_n < \epsilon/4. \quad (6)$$

Part 2: Set $N = \max\{Q_{n+2}(x), Q'(x), N'(x)\}$ and $\nu = \xi(x)$. Given $q \geq N$ we set $L = \max\{P, S'(x, q), S''(x, q), S_{n+2}(x, q)\}$. Combining (4) and (5), we have

$$d(f_q(x), z) < \nu \text{ and } s \geq L \Rightarrow |g_s(z)^* \bullet g_s(z) - g_s(f_q(x))^* \bullet g_s(f_q(x))|_n < \epsilon/2. \quad (7)$$

As $q \geq N'$, we have $d(f_q(x)^* \bullet f_q(x), 0) < \eta$, and putting this into (3) we get

$$s \geq L \Rightarrow |g_s(f_q(x)^* \bullet f_q(x))|_n < \epsilon/4. \quad (8)$$

Combining this with (6) gives

$$s \geq L \Rightarrow |g_s(f_q(x))^* \bullet g_s(f_q(x))|_n < \epsilon/2, \quad (9)$$

which together with (7) gives the result. $\square$

Lemma 8.2 *For all $x \in \mathcal{U}(A)$, all $n \in \mathbb{N}$ and all $\epsilon > 0$, there is an $\eta_n(x, \epsilon) > 0$ and a $T_n(x, \epsilon) \geq 0$ so that for all $y \in A$ and all $p \in [0, 1]$*

$$t \geq T_n \text{ and } d(y, x) < \eta_n \Rightarrow |\vartheta_B(p(f_t(y)^* \bullet f_t(y)))|_n < \epsilon.$$

Proof The function $\vartheta_B : V_B \rightarrow B$ is continuous, so there is a $m(\epsilon, n) \in \mathbb{N}$ and a $\delta(\epsilon, n)$ so that

$$\forall b \in A_{sad} \quad |b|_m < \delta \Rightarrow [b \in V_B \text{ and } |\vartheta_B(b)|_n < \epsilon].$$

The problem then reduces to finding conditions under which $|f_t(y)^* \bullet f_t(y)|_m < \delta$, and this is essentially contained in the proof of lemma 5.1. $\square$

Definition 8.3 For values $t \in [0, \infty)$, $u \in \mathcal{U}(A)$ and $p \in [0, 1]$ for which $f_t(u)^* \bullet f_t(u) \in V_B$ and $f_t(u) \bullet f_t(u)^* \in V_B$, we define $\hat{f}_{t,p}(u) = f_t(u) \bullet \vartheta(p(f_t(u)^* \bullet f_t(u)))$.

Lemma 8.4 The property $\Psi(x, y, s, t)$ defined as follows satisfies the conditions of lemma 2.1. For $x, y \in X = \mathcal{U}(A)$ and $s, t \geq 0$, $\Psi(x, y, s, t)$ is the logical value of the statement

$$\begin{aligned} & f_t(y)^* \bullet f_t(y) \in V_B \quad \text{and} \quad f_t(y) \bullet f_t(y)^* \in V_B \quad \text{and} \\ & \forall p \in [0, 1] \quad [g_s(\hat{f}_{t,p}(y))^* \bullet g_s(\hat{f}_{t,p}(y)) \in V_C \quad \text{and} \quad g_s(\hat{f}_{t,p}(y)) \bullet g_s(\hat{f}_{t,p}(y))^* \in V_C] \end{aligned} \quad (10)$$

Proof Part 1: By 5.1, for every $x \in \mathcal{U}(A)$ there is a $T(x) \geq 0$ and an open neighbourhood $O(x)$ of x in A so that

$$\forall t \geq T(x) \quad \forall y \in O(x) \quad f_t(y) \bullet f_t(y)^* \in V_B \quad \text{and} \quad f_t(y)^* \bullet f_t(y) \in V_B .$$

By lemma 8.1, given $x \in \mathcal{U}(A)$, there is an $N(x) \geq 0$ and a $\nu(x) > 0$ so that, for $q \geq N$ there is an $L(x, q) \geq 0$ so that

$$\forall z \in B \quad [d(f_q(x), z) < \nu \quad \text{and} \quad s \geq L] \Rightarrow [g_s(z)^* \bullet g_s(z) \in V_C \quad \text{and} \quad g_s(z) \bullet g_s(z)^* \in V_C] .$$

We take $q \geq \max\{T, N\} + 1$, $y \in O(x) \cap \mathcal{U}(A)$ and $|t - q| < 1$ in what follows, so $\hat{f}_{t,p}(y)$ (see 8.3) is defined for all $p \in [0, 1]$.

Part 2: Take $n \in \mathbb{N}$ and $1 > \chi > 0$ so that in B , $|z - f_q(x)|_n < \chi$ implies $d(f_q(x), z) < \nu$. Now consider the inequality

$$|f_t(y) \bullet b - f_q(x)|_n \leq |f_t(y) - f_q(x)|_{n+1} + |b|_{n+1} (1 + |f_q(x)|_{n+1} + |f_t(y) - f_q(x)|_{n+1}) , \quad (11)$$

where $b = \vartheta(p(f_t(y)^* \bullet f_t(y)))$, so that $f_t(y) \bullet b = \hat{f}_{t,p}(y)$. As f is pointwise bounded there is an $M(x, n+1)$ so that $|f_q(x)|_{n+1} \leq M$ for all $q \geq 0$. The conditions

$$|f_t(y) - f_q(x)|_{n+1} < \chi/2 \quad \text{and} \quad |\vartheta(p(f_t(y)^* \bullet f_t(y)))|_{n+1} < \chi/(4 + 2M) \quad (12)$$

and $s \geq L(x, q)$ then imply $\Psi(x, y, s, t)$.

Part 3: From 2.4 and 8.2 there are $T_{n+1}, \bar{P} \geq 0$ and $\eta_{n+1}, \bar{\eta} > 0$ (all depending on x) and, given $q \geq \bar{P}$, a $\bar{\delta} > 0$ (depending on x and q) so that the conditions in (12) are true if $t \geq T_{n+1}$, $|t - q| < \bar{\delta}$ and $d(x, y) < \min\{\eta_{n+1}, \bar{\eta}\}$. Now define $Q(x) = \max\{T_{n+1}, \bar{P}, T, N\} + 1$, and choose $\min\{\eta_{n+1}, \bar{\eta}\} > \delta(x) > 0$ so that $d(x, y) < \delta$ implies $y \in O(x)$. Given $q \geq Q$, set $R(x, q) = L(x, q)$ and $\epsilon(x, q) = \min\{\bar{\delta}, 1\}$. $\square$

Corollary 8.5 *There are continuous functions $\gamma : \mathcal{U} \rightarrow [0, \infty)$ and $\theta : [0, \infty) \rightarrow [0, \infty)$ (increasing) so that the formula*

$$r_{s,t,p}(u) = g_s(\hat{f}_{t,p}(u)) \bullet \vartheta_C(g_s(\hat{f}_{t,p}(u))^* \bullet g_s(\hat{f}_{t,p}(u)))$$

defines a continuous function

$$r : \{ (u, s, t, p) \in \mathcal{U}(A) \times [0, \infty) \times [0, \infty) \times [0, 1] : t \geq \gamma(u) \text{ and } s \geq \theta(t) \} \rightarrow \mathcal{U}(C) .$$

Proof Use 2.1 on 8.4. $\square$

Theorem 8.6 $\mathcal{U}(g \circ f) = \mathcal{U}(g) \circ \mathcal{U}(f) \in [\mathcal{U}(A), \mathcal{U}(C)]$.

Proof By 7.4 there is a valid reparameterisation $\phi : [0, \infty) \rightarrow [0, \infty)$ for g and f . Take the functions $\gamma : \mathcal{U}(A) \rightarrow [0, \infty)$ and $\theta : [0, \infty) \rightarrow [0, \infty)$ from lemma 8.5, and define $\psi : [0, \infty) \rightarrow [0, \infty)$ by $\psi(t) = \max\{t, \phi(t), \theta(t)\}$. As f , g and $g \circ_\psi f$ are PBAMs, we can generate respective functions $\alpha : \mathcal{U}(A) \rightarrow [0, \infty)$, $\beta : \mathcal{U}(B) \rightarrow [0, \infty)$ and $\mu : \mathcal{U}(A) \rightarrow [0, \infty)$ from 5.2. Define $\lambda : \mathcal{U}(A) \rightarrow [0, \infty)$ by $\lambda(u) = \max\{\alpha(u), \gamma(u)\}$ and $\omega : \mathcal{U}(A) \rightarrow [0, \infty)$ by $\omega(u) = \max\{\alpha(u), \beta(\tilde{f}_\lambda(u)), \gamma(u), \mu(u)\}$.

Now the function $h : \mathcal{U}(A) \times [0, 1] \rightarrow \mathcal{U}(C)$ defined by $h_p(u) = r_{\psi(\omega(u)), \omega(u), p}(u)$ gives a homotopy between $h_0(u) = (g \circ_\psi f)_\omega(u)$ and

$$h_1(u) = g_{\psi(\omega(u))}(\tilde{f}_\omega(u)) \bullet \vartheta_C(g_{\psi(\omega(u))}(\tilde{f}_\omega(u))^* \bullet g_{\psi(\omega(u))}(\tilde{f}_\omega(u))) .$$

By using $\tau, \zeta : [0, 1] \times \mathcal{U}(A) \rightarrow [0, \infty)$ defined by $\tau(p)(u) = (1 - p)\omega(u) + p\lambda(u)$ and $\zeta(p)(u) = \max\{\psi(\tau(p)(u)), \beta(\tilde{f}_\lambda(u))\}$ we define another homotopy $h' : \mathcal{U}(A) \times [0, 1] \rightarrow \mathcal{U}(C)$ by

$$h'_p(u) = g_{\zeta(p)(u)}(\tilde{f}_{\tau(p)}(u)) \bullet \vartheta_C(g_{\zeta(p)(u)}(\tilde{f}_{\tau(p)}(u))^* \bullet g_{\zeta(p)(u)}(\tilde{f}_{\tau(p)}(u))) .$$

This time $h'_0 = h_1$ as $\psi(\omega(u)) \geq \omega(u) \geq \beta(\tilde{f}_\lambda(u))$, and

$$h'_1(u) = g_{\zeta(1)(u)}(\tilde{f}_\lambda(u)) \bullet \vartheta_C(g_{\zeta(1)(u)}(\tilde{f}_\lambda(u))^* \bullet g_{\zeta(1)(u)}(\tilde{f}_\lambda(u))) .$$

Define another homotopy $h'' : \mathcal{U}(A) \times [0, 1] \rightarrow \mathcal{U}(C)$ by

$$h''_p(u) = g_{\eta(p)(u)}(\tilde{f}_\lambda(u)) \bullet \vartheta_C(g_{\eta(p)(u)}(\tilde{f}_\lambda(u))^* \bullet g_{\eta(p)(u)}(\tilde{f}_\lambda(u))) ,$$

where $\eta(p)(u) = (1 - p)\zeta(1)(u) + p\beta(\tilde{f}_\lambda(u))$. Now $h''_0 = h'_1$ and $h''_1 = \tilde{g}_\beta \circ \tilde{f}_\lambda$. $\square$

References

- [1] E.J. Beggs, *Strongly asymptotic morphisms on separable metrisable algebras*, Jour. of Functional Analysis **177**, 16-53 (2000).
- [2] A. Connes and N. Higson, *Déformations, morphismes asymptotiques et K -théorie bivariante*, C. R. Acad. Sci. Paris Sér. I Math. **311** no. 2 (1990), 101-106.
- [3] M. Dădărlat, *A note on asymptotic homomorphisms*, K -Theory **8** no. 5 (1994), 465-482.
- [4] T.W. Palmer, *$*$ -representations of U^* -algebras*, Indiana Univ. Math. J. **20** (1971), 929-933.
- [5] T.W. Palmer, *The Gelfand-Naimark pseudo norm on Banach $*$ -algebras*, J. Lond. Math. Soc. (2) **3** (1971), 59-66.
- [6] K. Thomsen, *Nonstable K -theory for operator algebras*, Internat. J. Math. **4** no. 3 (1991), 245-267.